

Citation for published version:

Chen, Y-C, Travers, RL, Walhin, J-P, Gonzalez, JT, Koumanov, F, Betts, JA & Thompson, D 2017, 'Feeding Influences Adipose Tissue Responses to Exercise in Overweight Men', American Journal of Physiology: Endocrinology and Metabolism, vol. 313, no. 1, pp. E84-E93. <https://doi.org/10.1152/ajpendo.00006.2017>

DOI:

[10.1152/ajpendo.00006.2017](https://doi.org/10.1152/ajpendo.00006.2017)

Publication date:

2017

Document Version

Peer reviewed version

[Link to publication](#)

University of Bath

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**FEEDING INFLUENCES ADIPOSE TISSUE RESPONSES TO EXERCISE IN
OVERWEIGHT MEN**

Running Head: Dietary status affects acute adipose responses to exercise

Yung-Chih Chen¹, Rebecca L. Travers¹, Jean-Philippe Walhin¹, Javier T. Gonzalez¹, Francoise
Koumanov¹, James A. Betts¹ and Dylan Thompson¹.

¹ Department for Health, University of Bath, United Kingdom

Corresponding author:

Professor Dylan Thompson

Department for Health

University of Bath

Bath, BA2 7AY

Email: d.thompson@bath.ac.uk

ABSTRACT

Feeding profoundly affects metabolic responses to exercise in various tissues but the effect of feeding status on human adipose tissue responses to exercise has never been studied. Ten healthy overweight men aged 26 ± 5 years (mean \pm SD) with a waist circumference of 105 ± 10 cm walked at 60% of maximum oxygen uptake under either FASTED or FED conditions in a randomised, counterbalanced design. Feeding comprised 648 ± 115 kcal 2 h before exercise. Blood samples were collected at regular intervals to examine changes in metabolic parameters and adipokine concentrations. Adipose tissue samples were obtained at baseline and one hour post-exercise to examine changes in adipose tissue mRNA expression and secretion of selected adipokines *ex-vivo*. Adipose tissue mRNA expression of PDK4, ATGL, HSL, FAT/CD36, GLUT4 and IRS2 in response to exercise were lower in FED compared to FASTED conditions (all $p \leq 0.05$). Post-exercise adipose IRS2 protein was affected by feeding ($p \leq 0.05$), but Akt2, AMPK, IRS1, GLUT4, PDK4 and HSL protein levels were not different. Feeding status did not impact serum and *ex-vivo* adipose secretion of IL-6, leptin or adiponectin in response to exercise. This is the first study to show that feeding prior to acute exercise affects post-exercise adipose tissue gene expression and we propose that feeding is likely to blunt long-term adipose tissue adaptation to regular exercise.

Keywords: Exercise, postprandial, adipose tissue, feeding

INTRODUCTION

It has become clear in the last 10 years or so that adipose tissue plays an active role in many physiological processes and pathological states (74, 78) and dysfunction within this tissue is characterised by tissue-specific insulin resistance, local inflammation, fibrosis, and the abnormal secretion of adipokines (36). Adipose tissue secretes dozens of mediators including the archetypal adipokines, adiponectin and leptin (16, 38, 46). Adiponectin is exclusively derived from adipose tissue and circulates in high concentrations (10–20 mg·L⁻¹). In other cases, the quantitative amounts secreted by adipose can be substantial, for example, subcutaneous adipose contributes up to a third of circulating interleukin-6 (IL-6) (45).

Feeding has a pronounced effect on the whole-body metabolic responses to exercise and reduces the contribution of fat towards metabolism (3, 23, 77). In addition, feeding influences the skeletal muscle responses to various forms of exercise (9, 10, 13). For example, pyruvate dehydrogenase kinase isozyme 4 (PDK4) is significantly up-regulated in muscle with exercise in fasted but not fed conditions (10). Other feeding-related changes in gene expression in muscle after exercise have been reported including altered expression of glucose transporter type 4 (GLUT4), PDK4, fatty acid translocase/CD36 (FAT/CD36), carnitine palmitoyltransferase 1 (CPT-1), uncoupling protein-3 (UCP3) and AMP-activated protein kinase (AMPK) (9). Collectively, therefore, there is strong evidence that feeding affects the responses to exercise in skeletal muscle. Adipose tissue plays a crucial role during exercise (18) and this led us to speculate that pre-exercise feeding may also affect the adipose tissue responses to acute exercise.

During moderate intensity exercise, adipose tissue provides much of the energy for working skeletal muscle through the mobilisation of stored triacylglycerol (27). Exercise also initiates

63 a number of other responses in adipose tissue such as increased blood flow and altered
64 expression of various adipokines within abdominal subcutaneous adipose tissue (66). It is
65 possible that these acute exercise-induced changes could be part of the mechanism through
66 which exercise improves health (66). However, all prior studies of adipose tissue responses to
67 exercise in humans have been conducted in the fasted state (8, 19, 25, 31, 32, 39). The effect
68 of feeding status on the response of human adipose tissue to exercise has never been studied.
69 This is despite the fact that feeding has a profound effect on adipose tissue (1, 30) and that we
70 spend the vast majority (~70%) of a 24-h period in a fed state (58). Consequently, the aim of
71 the present study was to investigate whether feeding influences the adipose tissue responses to
72 exercise.

METHODS

Ethical approval

The protocol was approved by Bristol Research Ethics Committee (REC reference number: 13/SW/0020) in accordance with the *declaration of Helsinki*. This trial is registered at ClinicalTrials.gov (ID: NCT02870075). All participants provided verbal and written informed consent before taking part.

Experimental design

Ten men aged 18 to 35 years with increased central adiposity were recruited via local advertisement. Participants attended the laboratory on three occasions for initial assessment of maximum oxygen uptake ($\dot{V}O_{2\max}$) and two subsequent main trials. The trial days involved walking for 60 min at 60% $\dot{V}O_{2\max}$ under either FASTED or FED conditions in a randomised, counterbalanced design separated by a 3–4 week wash-out period. This intensity and duration of exercise was selected because it relies heavily on fatty acids mobilised from adipose tissue and also because it is the type of exercise recommended in recent position stands (20, 27). Blood and adipose tissue were sampled at baseline and after exercise to examine the impact of prior feeding. There are no data regarding how different dietary status (fasted *versus* fed) affects the adipose responses to exercise. However, a previous study using a similar meal showed that feeding had an enormous effect on the use of lipid during exercise – and thus this indicates that the role of adipose tissue during exercise would be potentially very different (3, 77). Based on these results, in order to see an effect on lipid oxidation during exercise with 95% power and 5% alpha, we would require between 6–8 participants. We recruited ten men to account for greater variability in other outcome measures.

Inclusion and exclusion criteria

To be eligible to take part, participants were required to be overweight with a waist circumference of 94–128 cm (75). Participants were also required to be weight stable (63) for at least 3 months (mass stable $\pm 3\%$). Participants completed a health questionnaire to exclude any existing cardiovascular and metabolic diseases and a Physical Activity Readiness Questionnaire (PAR-Q) to make sure that participants were able to exercise safely. Individuals taking any medications known to influence lipid/carbohydrate metabolism or immune function and smokers were excluded. A summary of participants' physical characteristics is shown in Table 1.

Table 1. Participant physical characteristics (n = 10)

| Characteristics | Mean \pm SD |
|--|------------------|
| Age (years) | 26 \pm 5 |
| Body mass (kg) | 102.4 \pm 10.6 |
| Waist circumference (cm) | 105 \pm 10 |
| Hip circumference (cm) | 115 \pm 6 |
| Body mass index ($\text{kg}\cdot\text{m}^{-2}$) | 30.2 \pm 3.7 |
| Fat in L1-L4 region (kg) | 3.4 \pm 1.5 |
| $\dot{V}\text{O}_{2\text{max}}$ ($\text{mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$) | 42.4 \pm 6.4 |
| Systolic blood pressure (mmHg) | 132 \pm 21 |
| Diastolic blood pressure (mmHg) | 73 \pm 12 |

Fat in L1-L4 regions was determined using DEXA as previously described (22).

Pre-trial assessments

$\dot{V}O_{2max}$ measurement

An incremental continuous treadmill test until the point of volitional fatigue was used to determine $\dot{V}O_{2max}$. For most participants, a treadmill speed of 4 km·h⁻¹ and gradient of 8.5% was appropriate. This grade was maintained and the speed was increased by 1 km·h⁻¹ after every 3 min stage. One minute expired air samples were collected into Douglas bags (Hans Rudolph, MO, USA) and rate of perceived exertion (RPE) and heart rate were measured in the final minute of each stage and also at the point of volitional fatigue, defined as when the participant indicated that only 1 min remained until fatigue. Samples were analysed for relative expired fractions of oxygen and carbon dioxide (Servomex, Crowborough, UK) and the total volume within the bag was measured using a dry gas meter (Harvard Apparatus, Kent, UK).

Physical activity assessment

As part of the pre-trial assessments, participants wore a combined heart rate/accelerometer monitor for one week to assess their habitual physical activity energy expenditure (Actiheart, Cambridge Neurotechnology Ltd., Cambridge, UK). This was attached to the chest via 2 adhesive ECG pads for 24 h per day except for during showering/bathing/swimming (65).

Body composition analysis

Body mass was measured using digital scales following an overnight fast and post-void (TANITA corp., Tokyo, Japan). Waist and hip circumference was assessed according to World Health Organisation guidelines (75). Body composition was determined using Dual Energy X-ray Absorptiometry (DEXA; Discovery, Hologic, Bedford, UK). Abdominal subcutaneous and visceral adipose tissue mass was estimated from a central region between L1-L4 (22).

Trial days

In the 72 h prior to each main trial, participants were asked to refrain from performing any strenuous physical activity and from consuming alcohol/caffeine for 48 h prior to the main trials. A dietary record was completed 48 h before the first main trial and participants replicated this diet prior to their second main trial.

On main trial days, participants arrived at the laboratory between 8 and 9 am following a 12 h fast. After anthropometric measurements, participants rested on a bed for 15 min, followed by four 5-min expired gas sample collections to determine resting metabolic rate (RMR) (5, 11) using substrate oxidation under resting conditions (17). During exercise alternative equations were used (28).

After RMR assessment, a cannula was inserted into an antecubital forearm vein and a baseline venous blood sample was taken and allocated into tubes with either ethylenediaminetetraacetic acid (EDTA) or serum separation beads (Sarstedt Ltd, Leicester, UK). Plasma samples were centrifuged immediately at 3,465 g at 4 °C for 10 min. Serum samples were left for 45 min to coagulate before centrifugation. Subcutaneous adipose tissue samples (~1 g) were taken under local anaesthetic (1% lidocaine) from the area around the waist approximately 5 cm lateral to the umbilicus with a 14 G needle using an aspiration technique (72) followed by adipose tissue cleaning and processing as described previously (67).

Participants then either consumed a meal (FED) or remained fasting (FASTED) and cannula-derived venous blood samples were taken every 15 min for the following 60 min. A further blood sample was collected at 120 min immediately before the walking protocol. In both the

FASTED and FED treatments, participants walked on the treadmill at 60% $\dot{V}O_{2max}$ for 60 min and one minute of expired air samples, RPE and heart rate were collected at 5, 20, 40 and 60 min. After finishing exercise, another blood sample was immediately collected and participants then rested for a further 60 min. At this point, a second adipose tissue and final blood sample were taken.

Meal

The meal in the FED trial was the same as previously described in detail (7). The composition of the meal was selected to reflect typical breakfasts in the UK (7). The amount was normalised to resting energy requirements (7). Briefly, the total energy provided was 648 ± 115 kcal (carbohydrate 120.1 ± 21.3 g, fat 12.7 ± 2.3 g and protein 20.9 ± 3.7 g). The meal included white bread (Brace's thick white), cornflakes (Kellogg's cornflakes), semi-skimmed milk (Sainsbury; British semi skimmed milk), orange juice (Sainsbury; 100% pure squeezed smooth orange juice), spread (Unilever; I can't believe its not butter), jam (Sainsbury; strawberry jam) and sugar (Sainsbury; British white granulated sugar). Participants were asked to consume the meal within 15 min. In the FASTED trial, participants sat quietly for a 15-min period.

Adipose tissue gene expression and culture

After cleaning and mincing the adipose tissue biopsy sample, one portion of adipose tissue (approximately 200 mg) was immediately homogenised in 5 mL TRIzol (Invitrogen, Paisley, UK) in an RNase/DNase-free sterile tube (Invitrogen, Paisley, UK) and stored at -80°C before mRNA gene expression and protein analysis. The remaining adipose tissue was used for culture and four ~ 100 mg portions were placed in sterile culture plates (Nunc, Roskilde, Denmark) with endothelial cell basal media (ECBM) (Promocell, Germany) containing 0.1% fatty acid-free bovine serum albumin $100 \text{ U}\cdot\text{mL}^{-1}$ penicillin and $0.1 \text{ mg}\cdot\text{mL}^{-1}$ streptomycin (Sigma-

Aldrich, Gillingham, UK). Adipose tissue was incubated with a final ratio of 100 mg tissue per 1 mL ECBM media for 3 h (68) at 37 °C in a 5% CO₂ and 95 ± 5% relative humidity incubator (MCO-18A1C CO₂ incubator; Sanyo, Osaka, Japan). After the 3-h incubation, media was transferred to sterile tubes and stored at –80 °C. Adipokine secretion from adipose explants was normalised to explant adipose mass and then L1-L4 fat mass as described (67).

Real-time PCR

An RNeasy Mini Kit (Qiagen, Crawley, UK) was used to extract RNA from adipose tissue as described (72). Tissue samples were quantified using a Qubit 2.0 fluorimeter (Life Technologies, Paisley, UK). RNA was reversed transcribed (1 µg) to cDNA using a High Capacity Reverse Transcription Kit (Applied Biosystems, Warrington, UK). Organic phenol-chloroform phase from the RNA extraction was kept for further protein analysis. Real-time PCR was performed using a StepOne™ (Applied Biosystems, Warrington, UK). Predesigned primers and probes were obtained from Applied Biosystems for the measurement of expression of interleukin 6 (IL-6) (Hs00985639_m1), adiponectin (Hs00605917_m1), leptin (Hs00174877_m1), interleukin 18 (IL-18) (Hs00155517_m1), tumour necrosis factor alpha (TNF-α) (Hs99999043_m1), monocyte chemoattractant protein-1 (MCP-1) (Hs00234140_m1), 5' AMP-activated protein kinase (AMPK) (Hs01562315_m1 and Hs00178903_m1 combined), glucose transporter type 4 (GLUT4) (Hs00168966_m1), hormone-sensitive lipase (HSL) (Hs00193510_m1), insulin receptor substrate 1 (IRS1) (Hs00178563_m1), insulin receptor substrate 2 (IRS2) (Hs00275843_s1), sterol regulatory element binding protein 1c (SREBP-1c) (Hs01088691_m1), pyruvate dehydrogenase kinase isozyme (PDK4) (Hs00176875_m1), peroxisome proliferator-activated receptor γ (PPARγ) (Hs01115513_m1), peroxisome proliferator-activated receptor-gamma coactivator 1 alpha (PGC-1α) (Hs01016719_m1), RAC-alpha serine/threonine-protein kinase (Akt1) (Hs00178289_m1), adipose triglyceride

215 lipase (ATGL) (Hs00386101_m1), fatty acid translocase (FAT)/CD36 (Hs00169627_m1),
216 forkhead box protein O1 (FOXO1) (Hs01054576_m1), hexokinase 2 (HK2)
217 (Hs00606086_m1), PI3K-85 α (PIK3R1) (Hs00933163_m1), carnitine palmitoyltransferase 1B
218 (CPT1B) (Hs03046298_s1), The G0/G1 switch gene 2 (G0S2) (Hs00605971_m1),
219 peptidylpropyl isomerase A (PPIA) was used as an endogenous control (48). The comparative
220 threshold cycle (Ct) method was used to process data where $\Delta Ct = Ct \text{ target gene} - Ct \text{ PPIA}$.
221 Ct target genes were normalised to an internal calibrator (lowest ΔCt for each target gene) and
222 baseline. The Ct values for IL-6 (31 out of 40 samples), TNF- α (16 out of 40 samples), and IL-
223 18 (37 out of 40 samples) were frequently over 35 and thus these results are not included.

224

225 **Western blotting**

226 The adipose tissue protein fraction was isolated from the TRIzol phenol-chloroform phase
227 following the manufacturer's protocol (TRIzol Reagent, Life Technologies). Briefly, 1 mL of
228 organic phase was mixed with 1.5 mL isopropanol. After mixing, the samples were incubated
229 for 10 min at room temperature, followed by 10 min centrifugation at 12,000 g at 4 °C to pellet
230 the protein. One millilitre of protein pellet was washed using 2 mL of 0.3 M guanidine
231 hydrochloride in 95% ethanol for 20 min incubation followed by centrifugation at 7,500 g for
232 5 min at 4 °C. This process was repeated 3 times. After finishing the washing procedure, 2 mL
233 of 100% ethanol was added to the protein pellet for a further 20 min incubation at room
234 temperature before being centrifuged. The pellet was then left to air dry for 5–10 min. Then,
235 200 μL of 1% SDS was added to resuspended the pellet. The protein content of the samples
236 was determined using a BCA protein assay kit (Thermo Scientific, Waltham, USA). Proteins
237 (25 $\mu\text{g} \cdot \text{lane}^{-1}$) were separated by SDS-PAGE and transferred using a semidry electro-transfer
238 method to a nitrocellulose membrane. Western blotting analysis was performed with the
239 following antibodies: RAC-beta serine/threonine-protein kinase (Akt2)/PKB β (Millipore) (34),

AMPK (Cell Signalling Technology, USA) (72), GLUT4 (26), IRS1 (Millipore) (35), IRS2 (Millipore) (47), PDK4 (ABGENT, San Diego, USA) (29), HSL (Cell Signalling Technology, USA) (56) and glyceraldehyde 3-phosphate dehydrogenase (GAPDH) (Proteintech, USA) (62). The images were acquired in an EPI Chemi II darkroom (UVP) and bands quantified using VisionWorks LS analysis software (UVP).

Biochemical analysis

Plasma glucose and non-esterified fatty acids (NEFA) were measured using commercially available assay kits and analyser (Daytona Rx; Randox, Crumlin, UK). Serum insulin (Merckodia, Uppsala, Sweden) and both serum and adipose media concentrations of IL-6, leptin and adiponectin (R&D systems) were measured using Enzyme-linked immunosorbent assay (ELISA).

Statistics

Descriptive data are presented as means \pm standard deviation (SD). The variance bars on figures are presented as means with 95% confidence intervals (CI). Time series data were analysed using a two-way ANOVA (trial \times time) with repeated measures irrespective of minor deviations from a normality of distribution (43) using SPSS version 22 (IBM, Armonk, NY, USA). Where significant interactions (trial \times time) were found, post hoc paired *t*-tests were used to determine changes over time. Analysis of gene and protein expression data were conducted using logged transformed data as previously described (72). Incremental area under curve (iAUC) was calculated for insulin, glucose and NEFA using the trapezoid method (76) and analysed using paired *t*-tests. Statistical significance was set at $p \leq 0.05$.

RESULTS

Energy expenditure and substrate oxidation during FASTED and FED trials

There were modest feeding-induced differences between trials for relative exercise intensity (% $\dot{V}O_{2\max}$) and exercise energy expenditure (Table 2). Fat oxidation during exercise was reduced by ~45% in the FED trial (Table 2). Pre-fed resting metabolic rate (RMR) was not different ($2,103 \pm 418$ *versus* $2,058 \pm 365$ kcal·d⁻¹ in FASTED and FED trials, respectively).

Table 2. Physiological responses during 60 min exercise (n = 10).

| | FASTED | FED |
|---|-------------------|-----------------|
| Treadmill speed (km·h ⁻¹) | 5.7 ± 0.7 | 5.7 ± 0.7 |
| $\dot{V}O_{2\max}$ (%) | 59 ± 3 * | 60 ± 3 |
| Heart Rate (beat·min ⁻¹) | 155 ± 13 | 155 ± 14 |
| RPE (6–20) | 12 ± 2 | 12 ± 2 |
| Energy expenditure (kcal·h ⁻¹) | 746 ± 129 * | 771 ± 135 |
| Respiratory exchange ratio ($\dot{V}CO_2:\dot{V}O_2$) | 0.93 ± 0.03 * | 0.97 ± 0.03 |
| Carbohydrate oxidation (g·h ⁻¹) | 147 ± 41 * | 167 ± 39 |
| Fat oxidation (g·h ⁻¹) | 16 ± 8 * | 9 ± 6 |

Values are means \pm SD. * denotes significantly different between FASTED *versus* FED condition ($p \leq 0.05$).

Plasma glucose, NEFA and serum insulin concentrations

Trial \times Time interaction effects were found for blood glucose, insulin and NEFA between FASTED and FED trials ($p = 0.026$, $p = 0.003$ and $p = 0.001$, respectively). As expected, iAUCs for glucose and insulin were elevated for all parameters in the FED trial ($p = 0.02$ and $p = 0.03$, respectively) (Figure 1B and D). Peak glucose and insulin concentrations in the FED trial were reached 15 min post-meal (7.1 ± 0.6 mmol·L⁻¹ and 370 ± 185 pmol·L⁻¹, Figure 1A

and C, respectively). NEFA concentrations were lower at most time points in the FED trial (Figure 1E).

[INSERT FIGURE 1 ABOUT HERE]

Adipose tissue mRNA expression

There was an interaction effect for adipose tissue PDK4, ATGL, HSL, FAT/CD36, GLUT4 and IRS2 (all $p \leq 0.05$) after exercising under FASTED versus FED conditions (Figure 2). These interaction effects are explained by divergent responses between FASTED and FED trials with a larger increase in FASTED conditions when compared to either a smaller change or a modest decrease in FED conditions (Figure 2). There was a time effect for HK2, MCP-1 and PGC-1 α (Figure 2). The expression of the remaining genes was not significantly different between trials or over time (Figure 2).

[INSERT FIGURE 2 ABOUT HERE]

Adipose tissue protein expression

There was an interaction effect for IRS2 protein expression between trials ($p \leq 0.05$), showing approximately a 2-fold increase in FASTED conditions, and no change in FED conditions (Figure 3A). The change in the expression of the remaining proteins was not statistically different between FASTED and FED conditions. Representative Western blots are shown in Figure 3B.

[INSERT FIGURE 3 ABOUT HERE]

Serum adipokine concentrations and adipose tissue secretion *ex vivo*

Serum IL-6 increased (Figure 4A) and serum leptin decreased (Figure 4C) in both trials ($p < 0.05$). There was a modest time effect for serum adiponectin (Figure 4E), although no interaction effects were identified for serum IL-6, leptin and adiponectin. Furthermore, no time or time \times trial interactions were identified for *ex vivo* adipose explant secretion of IL-6, leptin and adiponectin in response to exercise under FASTED and FED conditions (Figure 4).

[INSERT FIGURE 4 ABOUT HERE]

DISCUSSION

This study presents the first evidence that feeding status alters the human adipose tissue response to acute exercise and thus feeding status has the potential to influence the long-term adaptation of adipose tissue to regular exercise.

In the present study, our feeding protocol successfully manipulated systemic concentrations of glucose and insulin. In the two hours prior to exercise, there was a 168-fold and 26-fold difference in insulin and glucose iAUC, respectively. As anticipated with this study design, prior feeding increased relative carbohydrate utilisation and decreased fat oxidation during exercise (3, 23, 77). Thus, exercise in FED and FASTED trials was performed in a very different physiological state.

At the gene expression level, we found that adipose tissue responded differently to moderate-intensity exercise under FASTED *versus* FED conditions. When compared to the changes in fasted exercise conditions, feeding led to lower changes or a decrease in PDK4, ATGL, HSL, FAT/CD36, GLUT4 and IRS2 mRNA as demonstrated by interaction effects for these outcomes. Over time, acute differences in skeletal muscle gene expression with exercise conducted in the fasted *versus* the fed state have been proposed to contribute to diverse physiological adaptations (12, 70). Our data demonstrate that feeding status also alters adipose tissue responses to an acute bout of exercise. We have previously shown that weight loss leads to a large increase in genes such as PDK4 and HSL in adipose tissue (71) whereas overfeeding leads to a profound decrease in both PDK4 and HSL in adipose tissue (72). Thus, feeding has the potential to affect the acute adipose tissue responses to exercise and, given the important role of adipose tissue in health and the nature of these changes (66, 71, 72), we propose that feeding before exercise blunts some of the health-related changes induced by exercise training.

We found a difference in adipose IRS2 protein content between FASTED and FED exercise conditions although the other measured proteins were not affected. We should highlight that our protein measurements represent only total protein content. Repeated small changes in adipose protein synthesis are likely lead to accumulated differences and functional changes in adipose phenotype over time (6, 50). Clearly, the only way to know if the acute changes observed in the current study translate into long term differences in protein content in adipose is to examine whether chronic training conducted in the fasted *versus* postprandial state leads to divergent adaptations.

Carbohydrate and lipid metabolism in adipose tissue

A primary function of PDK4 is to regulate glucose metabolism by inhibiting pyruvate dehydrogenase complex activity. Fasting and exercise increase PDK4 mRNA expression in skeletal muscle (52, 53) and insulin suppresses PDK4 mRNA and protein content in skeletal muscle (37, 40). It is possible that insulin could be responsible for the lower adipose PDK4 mRNA response in the FED trial, although higher NEFA in the FASTED trial could also have increased PDK4 expression (2, 33). Other feeding-related studies have shown that muscle PDK4 mRNA expression remained unchanged 1–4 h after exercise in the fed state (with lower NEFA concentrations), when compared with exercise in the fasted state (9, 10). Thus, exercise in the fasted state appears to increase PDK4 expression in both muscle and adipose, whereas exercise in the fed state does not.

ATGL and HSL mobilise stored fat and release it into the circulation (79). In the current study, gene expression of ATGL, HSL and FAT/CD36 in adipose were all differentially expressed in FASTED and FED exercise conditions. These interaction effects are explained by responses in

the FED trial being either lower or in the opposite direction to the FASTED trial. There was also greater fat oxidation during exercise in the FASTED trial. Other studies have also shown similar responses in fasted exercise with higher fat oxidation and increased skeletal muscle FAT/CD36 gene expression (9). Moreover, these findings are consistent with the previously observed increase in adipose HSL activity reported during cycling exercise, which is blunted with nicotinic acid ingestion (73). Thus, given the nature and direction of these changes, we propose that feeding blunts at least some of the exercise-induced stimulus on adipose tissue.

We found GLUT4 mRNA, IRS2 mRNA and IRS2 protein were also differentially expressed in adipose in FASTED and FED exercise trials. These effects were subtle but consistent, and this also seems to indicate that exercise in a fed state will not generate the same change in pathways involved in glucose metabolism and signalling within adipose tissue as fasted exercise. Previous findings in skeletal muscle have found an increase in GLUT4 mRNA in fasted but not fed conditions after exercise (9, 10).

PGC-1 α mRNA is a transcriptional coactivator involved in mitochondrial biogenesis (60). Acute exercise increases PGC-1 α mRNA expression in human skeletal muscle (21, 54) and in rodent white adipose tissue (64). However, the impact of feeding status and/or carbohydrate variability prior to exercise on skeletal muscle PGC-1 α mRNA is controversial. Some studies indicate that PGC-1 α mRNA expression is up-regulated post-exercise whether acute exercise is performed in fed or fasted conditions (10, 42) but other studies show that higher carbohydrate availability prior to exercise blunts PGC1- α mRNA expression in skeletal muscle both at rest and post-exercise (4, 55). In the present study, the increase in adipose PGC-1 α mRNA expression after exercise was unaffected by feeding status. Chronic training studies indicate that adipose PGC1- α mRNA is increased in humans (59) and rodents (64, 69) and this has also

been reported to increase PGC1- α protein content and mitochondrial biogenesis in rodents (69). Whether the present results indicate an acute exercise-induced increase in mitochondrial biogenesis in human adipose tissue is plausible but unclear at the present time.

Adipokines response to exercise and impact of feeding status

Previous studies have shown that circulating adipokine concentrations are affected by acute exercise (8) and energy consumption during exercise alters these systemic responses (57, 61). However, evidence from studies that have manipulated feeding status *prior to* exercise is scarce. Zoladz et al. (80) found no difference in circulating IL-6 and leptin after a single bout of exercise, in a fed or fasted state. However, the duration of exercise lasted only 12 min and this might be insufficient to examine the notion that feeding status influences circulating adipokines. The exercise in the present study was 60 min, and we too found no evidence that pre-exercise feeding affects circulating IL-6, leptin and adiponectin. This may be due to the fact that moderate intensity exercise has only a modest effect on many of these parameters and thus there is little potential for feeding to interact with exercise and exert an effect (41). Serum IL-6 was increased over time in both fasted and fed trials in the present study, which could be partly due to the effect of exercise (51). As we did not observe changes of adipose IL-6 secretion *ex-vivo*, the increase in serum IL-6 might be caused by release from skeletal muscle during exercise (49), but it is also possible that reflects local production of IL-6 due to prolonged cannulation (14).

Temporal and population-specific considerations

This study is the first to examine the impact of feeding on adipose tissue responses to exercise. We recruited overweight participants and this focus is a strength given that ~62% of the UK population are overweight (24). Increased adiposity has a profound effect on adipose tissue

function (e.g., a down regulation of GLUT 4 mRNA (67) and reduction in postprandial adipose tissue blood flow (44)). These could be important considerations when interpreting our findings. We should also highlight that in the current study we were limited to only two adipose tissue biopsies due to a concern over potential interference from repeated sampling (15) and so we do not have a full and complete time course. For the first study of this kind and with limited sampling opportunities, a second biopsy 60 min post-exercise was considered to balance the requirement to capture pathways that rapidly change and those that are slower to respond. Subsequent studies should consider more frequent adipose sampling and/or the inclusion of additional resting trial(s). Depending on the kinetics of each response, our sample timing framework will be appropriate for some outcomes and less appropriate for others. Furthermore, in the absence of an adipose biopsy immediately prior to exercise, it is hard to establish whether some effects in the FASTED trial are due to the modestly extended fasting period or due to exercise (or the interaction between fasting and exercise). What is very apparent, however, is that exercise in FED conditions does not lead to the same changes as exercise in FASTED conditions.

Conclusion

This study provides the first evidence that the feeding status alters the response of adipose tissue to acute exercise. Several genes involved in lipid metabolism, insulin signaling and glucose transport were differentially expressed in adipose tissue when exercise was performed in a fed *versus* fasted state with either lower or opposing responses after feeding. Given the nature and direction of these differences, we propose that feeding is likely to blunt long-term adaptations induced within adipose tissue in response to regular exercise.

ACKNOWLEDGMENTS

We thank all the participants for their time and effort to take part in this project.

GRANTS

This project was funded in part by the Ministry of Education (Taiwan) and the University of Bath. Thompson and Travers were funded by a grant from BBSRC (BB/N004809/1) and Koumanov was funded by a grant from MRC (MR/P002927/1).

DISCLOSURES

The authors declare no competing interests.

AUTHOR CONTRIBUTIONS

Yung-Chih Chen was responsible for study design and conduct, data collection, data analysis, data interpretation, statistical analysis, and manuscript revision; Rebecca L. Travers was responsible for mRNA gene expression data collection and manuscript revision. Jean-Philippe Walhin was responsible for mRNA gene expression analysis and manuscript revision. Javier T. Gonzalez was responsible for manuscript revision. Francoise Koumanov was responsible for western blotting analysis, interpretation, and manuscript revision. James Betts was responsible for study design, and manuscript revision; Dylan Thompson was responsible for funding, study design, data interpretation, and manuscript revision.

459 REFERENCES

- 460 1. **Alligier M, Meugnier E, Debard C, Lambert-Porcheron S, Chanseau E, Sothier M, Loizon E,**
461 **Hssain AA, Brozek J, Scoazec JY, Morio B, Vidal H, and Laville M.** Subcutaneous adipose tissue
462 remodeling during the initial phase of weight gain induced by overfeeding in humans. *The Journal of*
463 *clinical endocrinology and metabolism* 97: E183-192, 2012.
- 464 2. **Arkininstall MJ, Tunstall RJ, Cameron-Smith D, and Hawley JA.** Regulation of metabolic genes
465 in human skeletal muscle by short-term exercise and diet manipulation. *American journal of*
466 *physiology Endocrinology and metabolism* 287: E25-31, 2004.
- 467 3. **Backhouse SH, Williams C, Stevenson E, and Nute M.** Effects of the glycemic index of
468 breakfast on metabolic responses to brisk walking in females. *European journal of clinical nutrition* 61:
469 590-596, 2007.
- 470 4. **Bartlett JD, Louhelainen J, Iqbal Z, Cochran AJ, Gibala MJ, Gregson W, Close GL, Drust B, and**
471 **Morton JP.** Reduced carbohydrate availability enhances exercise-induced p53 signaling in human
472 skeletal muscle: implications for mitochondrial biogenesis. *American journal of physiology Regulatory,*
473 *integrative and comparative physiology* 304: R450-458, 2013.
- 474 5. **Betts JA, Thompson D, Richardson JD, Chowdhury EA, Jeans M, Holman GD, and Tsintzas K.**
475 Bath Breakfast Project (BBP)--examining the role of extended daily fasting in human energy balance
476 and associated health outcomes: study protocol for a randomised controlled trial [ISRCTN31521726].
477 *Trials* 12: 172, 2011.
- 478 6. **Booth FW and Neufer PD.** Exercise controls gene expression. *Am Sci* 93: 28-35, 2005.
- 479 7. **Chowdhury EA, Richardson JD, Tsintzas K, Thompson D, and Betts JA.** Carbohydrate-rich
480 breakfast attenuates glycaemic, insulinaemic and ghrelin response to ad libitum lunch relative to
481 morning fasting in lean adults. *The British journal of nutrition* 114: 98-107, 2015.
- 482 8. **Christiansen T, Bruun JM, Paulsen SK, Olholm J, Overgaard K, Pedersen SB, and Richelsen B.**
483 Acute exercise increases circulating inflammatory markers in overweight and obese compared with
484 lean subjects. *European journal of applied physiology* 113: 1635-1642, 2013.
- 485 9. **Civitarese AE, Hesselink MK, Russell AP, Ravussin E, and Schrauwen P.** Glucose ingestion
486 during exercise blunts exercise-induced gene expression of skeletal muscle fat oxidative genes.
487 *American journal of physiology Endocrinology and metabolism* 289: E1023-1029, 2005.
- 488 10. **Cluberton LJ, McGee SL, Murphy RM, and Hargreaves M.** Effect of carbohydrate ingestion on
489 exercise-induced alterations in metabolic gene expression. *Journal of applied physiology* 99: 1359-
490 1363, 2005.
- 491 11. **Compher C, Frankenfield D, Keim N, Roth-Yousey L, and Evidence Analysis Working G.** Best
492 practice methods to apply to measurement of resting metabolic rate in adults: a systematic review.
493 *Journal of the American Dietetic Association* 106: 881-903, 2006.
- 494 12. **De Bock K, Derave W, Eijnde BO, Hesselink MK, Koninckx E, Rose AJ, Schrauwen P, Bonen A,**
495 **Richter EA, and Hespel P.** Effect of training in the fasted state on metabolic responses during exercise
496 with carbohydrate intake. *Journal of applied physiology* 104: 1045-1055, 2008.
- 497 13. **De Bock K, Richter EA, Russell AP, Eijnde BO, Derave W, Ramaekers M, Koninckx E, Leger B,**
498 **Verhaeghe J, and Hespel P.** Exercise in the fasted state facilitates fibre type-specific intramyocellular
499 lipid breakdown and stimulates glycogen resynthesis in humans. *The Journal of physiology* 564: 649-
500 660, 2005.
- 501 14. **Dixon NC, Hurst TL, Talbot DC, Tyrrell RM, and Thompson D.** Active middle-aged men have
502 lower fasting inflammatory markers but the postprandial inflammatory response is minimal and
503 unaffected by physical activity status. *Journal of applied physiology* 107: 63-68, 2009.
- 504 15. **Dordevic AL, Pendergast FJ, Morgan H, Villas-Boas S, Caldow MK, Larsen AE, Sinclair AJ, and**
505 **Cameron-Smith D.** Postprandial Responses to Lipid and Carbohydrate Ingestion in Repeated
506 Subcutaneous Adipose Tissue Biopsies in Healthy Adults. *Nutrients* 7: 5347-5361, 2015.

16. **Fain JN, Madan AK, Hiler ML, Cheema P, and Bahouth SW.** Comparison of the Release of Adipokines by Adipose Tissue, Adipose Tissue Matrix, and Adipocytes from Visceral and Subcutaneous Abdominal Adipose Tissues of Obese Humans. *Endocrinology* 145: 2273-2282, 2004.
17. **Frayn KN.** Calculation of substrate oxidation rates in vivo from gaseous exchange. *Journal of applied physiology: respiratory, environmental and exercise physiology* 55: 628-634, 1983.
18. **Frayn KN.** Fat as a fuel: emerging understanding of the adipose tissue-skeletal muscle axis. *Acta physiologica* 199: 509-518, 2010.
19. **Frydelund-Larsen L, Akerstrom T, Nielsen S, Keller P, Keller C, and Pedersen BK.** Visfatin mRNA expression in human subcutaneous adipose tissue is regulated by exercise. *American journal of physiology Endocrinology and metabolism* 292: E24-31, 2007.
20. **Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, Nieman DC, Swain DP, and American College of Sports M.** American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Medicine and science in sports and exercise* 43: 1334-1359, 2011.
21. **Gidlund EK, Ydfors M, Appel S, Rundqvist H, Sundberg CJ, and Norrbom J.** Rapidly elevated levels of PGC-1alpha-b protein in human skeletal muscle after exercise: exploring regulatory factors in a randomized controlled trial. *Journal of applied physiology* 119: 374-384, 2015.
22. **Glickman SG, Marn CS, Supiano MA, and Dengel DR.** Validity and reliability of dual-energy X-ray absorptiometry for the assessment of abdominal adiposity. *Journal of applied physiology* 97: 509-514, 2004.
23. **Gonzalez JT, Veasey RC, Rumbold PL, and Stevenson EJ.** Breakfast and exercise contingently affect postprandial metabolism and energy balance in physically active males. *The British journal of nutrition* 110: 721-732, 2013.
24. **Health and Social Care Information Centre.** Health Survey for England 2012 Trend Tables: Adult Trend Tables. on-line: NHS Digital, 2012.
25. **Hojbjerre L, Rosenzweig M, Dela F, Bruun JM, and Stallknecht B.** Acute exercise increases adipose tissue interstitial adiponectin concentration in healthy overweight and lean subjects. *European journal of endocrinology / European Federation of Endocrine Societies* 157: 613-623, 2007.
26. **Holman GD, Kozka IJ, Clark AE, Flower CJ, Saltis J, Habberfield AD, Simpson IA, and Cushman SW.** Cell surface labeling of glucose transporter isoform GLUT4 by bis-mannose photolabel. Correlation with stimulation of glucose transport in rat adipose cells by insulin and phorbol ester. *The Journal of biological chemistry* 265: 18172-18179, 1990.
27. **Horowitz JF.** Fatty acid mobilization from adipose tissue during exercise. *Trends in endocrinology and metabolism: TEM* 14: 386-392, 2003.
28. **Jeukendrup AE and Wallis GA.** Measurement of substrate oxidation during exercise by means of gas exchange measurements. *International journal of sports medicine* 26 Suppl 1: S28-37, 2005.
29. **Jiang LQ, de Castro Barbosa T, Massart J, Deshmukh AS, Lofgren L, Duque-Guimaraes DE, Ozilgen A, Osler ME, Chibalin AV, and Zierath JR.** Diacylglycerol kinase-delta regulates AMPK signaling, lipid metabolism, and skeletal muscle energetics. *American journal of physiology Endocrinology and metabolism* 310: E51-60, 2016.
30. **Johannsen DL, Tchoukalova Y, Tam CS, Covington JD, Xie W, Schwarz JM, Bajpeyi S, and Ravussin E.** Effect of 8 weeks of overfeeding on ectopic fat deposition and insulin sensitivity: testing the "adipose tissue expandability" hypothesis. *Diabetes care* 37: 2789-2797, 2014.
31. **Keller C, Keller P, Marshal S, and Pedersen BK.** IL-6 gene expression in human adipose tissue in response to exercise--effect of carbohydrate ingestion. *The Journal of physiology* 550: 927-931, 2003.
32. **Keller P, Keller C, Steensberg A, Robinson LE, and Pedersen BK.** Leptin gene expression and systemic levels in healthy men: effect of exercise, carbohydrate, interleukin-6, and epinephrine. *Journal of applied physiology* 98: 1805-1812, 2005.

33. **Kiilerich K, Gudmundsson M, Birk JB, Lundby C, Taudorf S, Plomgaard P, Saltin B, Pedersen PA, Wojtaszewski JF, and Pilegaard H.** Low muscle glycogen and elevated plasma free fatty acid modify but do not prevent exercise-induced PDH activation in human skeletal muscle. *Diabetes* 59: 26-32, 2010.
34. **Koumanov F, Pereira VJ, Richardson JD, Sargent SL, Fazakerley DJ, and Holman GD.** Insulin regulates Rab3-Noc2 complex dissociation to promote GLUT4 translocation in rat adipocytes. *Diabetologia* 58: 1877-1886, 2015.
35. **Lamphere L and Lienhard GE.** Components of signaling pathways for insulin and insulin-like growth factor-I in muscle myoblasts and myotubes. *Endocrinology* 131: 2196-2202, 1992.
36. **Langin D, Frühbeck G, Frayn KN, and Lafontan M.** Adipose tissue: development, anatomy and functions. In: *Obesity: Science to Practice*, edited by Williams G and Frühbeck G. Chichester, UK: Wiley-Blackwell, 2009, p. 79-108.
37. **Lee FN, Zhang L, Zheng D, Choi WS, and Youn JH.** Insulin suppresses PDK-4 expression in skeletal muscle independently of plasma FFA. *American journal of physiology Endocrinology and metabolism* 287: E69-74, 2004.
38. **Lehr S, Hartwig S, Lamers D, Famulla S, Müller S, Hanisch F-G, Cuvelier C, Ruige J, Eckardt K, Ouwens DM, Sell H, and Eckel J.** Identification and Validation of Novel Adipokines Released from Primary Human Adipocytes. *Molecular & Cellular Proteomics* 11, 2012.
39. **Leick L, Lindegaard B, Stensvold D, Plomgaard P, Saltin B, and Pilegaard H.** Adipose tissue interleukin-18 mRNA and plasma interleukin-18: effect of obesity and exercise. *Obesity* 15: 356-363, 2007.
40. **Majer M, Popov KM, Harris RA, Bogardus C, and Prochazka M.** Insulin downregulates pyruvate dehydrogenase kinase (PDK) mRNA: potential mechanism contributing to increased lipid oxidation in insulin-resistant subjects. *Molecular genetics and metabolism* 65: 181-186, 1998.
41. **Markovitch D, Tyrrell RM, and Thompson D.** Acute moderate-intensity exercise in middle-aged men has neither an anti- nor proinflammatory effect. *Journal of applied physiology* 105: 260-265, 2008.
42. **Mathai AS, Bonen A, Benton CR, Robinson DL, and Graham TE.** Rapid exercise-induced changes in PGC-1 α mRNA and protein in human skeletal muscle. *Journal of applied physiology* 105: 1098-1105, 2008.
43. **Maxwell SE and Delaney HD.** *Designing Experiments and Analyzing Data: A Model Comparison Perspective*. Belmont, CA, USA.: Wadsworth, 1990.
44. **McQuaid SE, Hodson L, Neville MJ, Dennis AL, Cheeseman J, Humphreys SM, Ruge T, Gilbert M, Fielding BA, Frayn KN, and Karpe F.** Downregulation of adipose tissue fatty acid trafficking in obesity: a driver for ectopic fat deposition? *Diabetes* 60: 47-55, 2011.
45. **Mohamed-Ali V, Goodrick S, Rawesh A, Katz DR, Miles JM, Yudkin JS, Klein S, and Coppack SW.** Subcutaneous adipose tissue releases interleukin-6, but not tumor necrosis factor- α , in vivo. *Journal of Clinical Endocrinology & Metabolism* 82: 4196-4200, 1997.
46. **Moro C, Klimcakova E, Lolmede K, Berlan M, Lafontan M, Stich V, Bouloumie A, Galitzky J, Arner P, and Langin D.** Atrial natriuretic peptide inhibits the production of adipokines and cytokines linked to inflammation and insulin resistance in human subcutaneous adipose tissue. *Diabetologia* 50: 1038-1047, 2007.
47. **Neukamm SS, Ott J, Dammeier S, Lehmann R, Haring HU, Schleicher E, and Weigert C.** Phosphorylation of serine 1137/1138 of mouse insulin receptor substrate (IRS) 2 regulates cAMP-dependent binding to 14-3-3 proteins and IRS2 protein degradation. *The Journal of biological chemistry* 288: 16403-16415, 2013.
48. **Neville MJ, Collins JM, Gloyd AL, McCarthy MI, and Karpe F.** Comprehensive human adipose tissue mRNA and microRNA endogenous control selection for quantitative real-time-PCR normalization. *Obesity* 19: 888-892, 2011.
49. **Pedersen BK.** Muscular interleukin-6 and its role as an energy sensor. *Medicine and science in sports and exercise* 44: 392-396, 2012.

50. **Perry CG, Lally J, Holloway GP, Heigenhauser GJ, Bonen A, and Spriet LL.** Repeated transient mRNA bursts precede increases in transcriptional and mitochondrial proteins during training in human skeletal muscle. *The Journal of physiology* 588: 4795-4810, 2010.
51. **Petersen AM and Pedersen BK.** The anti-inflammatory effect of exercise. *Journal of applied physiology* 98: 1154-1162, 2005.
52. **Pilegaard H, Ordway GA, Saltin B, and Neufer PD.** Transcriptional regulation of gene expression in human skeletal muscle during recovery from exercise. *American journal of physiology Endocrinology and metabolism* 279: E806-814, 2000.
53. **Pilegaard H, Saltin B, and Neufer PD.** Effect of short-term fasting and refeeding on transcriptional regulation of metabolic genes in human skeletal muscle. *Diabetes* 52: 657-662, 2003.
54. **Pilegaard H, Saltin B, and Neufer PD.** Exercise induces transient transcriptional activation of the PGC-1 α gene in human skeletal muscle. *The Journal of physiology* 546: 851-858, 2003.
55. **Psilander N, Frank P, Flockhart M, and Sahlin K.** Exercise with low glycogen increases PGC-1 α gene expression in human skeletal muscle. *European journal of applied physiology* 113: 951-963, 2013.
56. **Rittig N, Bach E, Thomsen HH, Pedersen SB, Nielsen TS, Jorgensen JO, Jessen N, and Moller N.** Regulation of Lipolysis and Adipose Tissue Signaling during Acute Endotoxin-Induced Inflammation: A Human Randomized Crossover Trial. *PloS one* 11: e0162167, 2016.
57. **Robson-Ansley P, Barwood M, Eglin C, and Ansley L.** The effect of carbohydrate ingestion on the interleukin-6 response to a 90-minute run time trial. *International journal of sports physiology and performance* 4: 186-194, 2009.
58. **Ruge T, Hodson L, Cheeseman J, Dennis AL, Fielding BA, Humphreys SM, Frayn KN, and Karpe F.** Fasted to fed trafficking of Fatty acids in human adipose tissue reveals a novel regulatory step for enhanced fat storage. *The Journal of clinical endocrinology and metabolism* 94: 1781-1788, 2009.
59. **Ruschke K, Fishbein L, Dietrich A, Kloting N, Tonjes A, Oberbach A, Fasshauer M, Jenkner J, Schon MR, Stumvoll M, Bluher M, and Mantzoros CS.** Gene expression of PPAR γ and PGC-1 α in human omental and subcutaneous adipose tissues is related to insulin resistance markers and mediates beneficial effects of physical training. *European journal of endocrinology / European Federation of Endocrine Societies* 162: 515-523, 2010.
60. **Scarpulla RC, Vega RB, and Kelly DP.** Transcriptional integration of mitochondrial biogenesis. *Trends in endocrinology and metabolism: TEM* 23: 459-466, 2012.
61. **Starkie RL, Arkinstall MJ, Koukoulas I, Hawley JA, and Febbraio MA.** Carbohydrate ingestion attenuates the increase in plasma interleukin-6, but not skeletal muscle interleukin-6 mRNA, during exercise in humans. *The Journal of physiology* 533: 585-591, 2001.
62. **Steen HC, Nogusa S, Thapa RJ, Basagoudanavar SH, Gill AL, Merali S, Barrero CA, Balachandran S, and Gamero AM.** Identification of STAT2 serine 287 as a novel regulatory phosphorylation site in type I interferon-induced cellular responses. *The Journal of biological chemistry* 288: 747-758, 2013.
63. **Stevens J, Truesdale KP, McClain JE, and Cai J.** The definition of weight maintenance. *International journal of obesity* 30: 391-399, 2006.
64. **Sutherland LN, Bomhof MR, Capozzi LC, Basaraba SA, and Wright DC.** Exercise and adrenaline increase PGC-1 α mRNA expression in rat adipose tissue. *The Journal of physiology* 587: 1607-1617, 2009.
65. **Thompson D, Batterham AM, Bock S, Robson C, and Stokes K.** Assessment of low-to-moderate intensity physical activity thermogenesis in young adults using synchronized heart rate and accelerometry with branched-equation modeling. *The Journal of nutrition* 136: 1037-1042, 2006.
66. **Thompson D, Karpe F, Lafontan M, and Frayn K.** Physical activity and exercise in the regulation of human adipose tissue physiology. *Physiological reviews* 92: 157-191, 2012.
67. **Travers RL, Motta AC, Betts JA, Bouloumie A, and Thompson D.** The impact of adiposity on adipose tissue-resident lymphocyte activation in humans. *International journal of obesity* 39: 762-769, 2015.

68. **Travers RL, Motta AC, Betts JA, and Thompson D.** Adipose tissue metabolic and inflammatory responses to a mixed meal in lean, overweight and obese men. *European journal of nutrition*, 2015.
69. **Trevellin E, Scorzeto M, Olivieri M, Granzotto M, Valerio A, Tedesco L, Fabris R, Serra R, Quarta M, Reggiani C, Nisoli E, and Vettor R.** Exercise training induces mitochondrial biogenesis and glucose uptake in subcutaneous adipose tissue through eNOS-dependent mechanisms. *Diabetes* 63: 2800-2811, 2014.
70. **Van Proeyen K, Szlufcik K, Nielens H, Ramaekers M, and Hespel P.** Beneficial metabolic adaptations due to endurance exercise training in the fasted state. *Journal of applied physiology* 110: 236-245, 2011.
71. **Walhin JP, Dixon NC, Betts JA, and Thompson D.** The impact of exercise intensity on whole body and adipose tissue metabolism during energy restriction in sedentary overweight men and postmenopausal women. *Physiological reports* 4, 2016.
72. **Walhin JP, Richardson JD, Betts JA, and Thompson D.** Exercise counteracts the effects of short-term overfeeding and reduced physical activity independent of energy imbalance in healthy young men. *The Journal of physiology* 591: 6231-6243, 2013.
73. **Watt MJ, Holmes AG, Steinberg GR, Mesa JL, Kemp BE, and Febbraio MA.** Reduced plasma FFA availability increases net triacylglycerol degradation, but not GPAT or HSL activity, in human skeletal muscle. *American journal of physiology Endocrinology and metabolism* 287: E120-127, 2004.
74. **Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, and Ferrante AW, Jr.** Obesity is associated with macrophage accumulation in adipose tissue. *The Journal of clinical investigation* 112: 1796-1808, 2003.
75. **WHO.** World Health Organisation-Waist Circumference and Waist-Hip Ratio: Report of a WHO Expert Consultation, 2008.
76. **Wolever TM and Jenkins DJ.** The use of the glycemic index in predicting the blood glucose response to mixed meals. *The American journal of clinical nutrition* 43: 167-172, 1986.
77. **Wu CL, Nicholas C, Williams C, Took A, and Hardy L.** The influence of high-carbohydrate meals with different glycaemic indices on substrate utilisation during subsequent exercise. *The British journal of nutrition* 90: 1049-1056, 2003.
78. **Xu HY, Barnes GT, Yang Q, Tan Q, Yang DS, Chou CJ, Sole J, Nichols A, Ross JS, Tartaglia LA, and Chen H.** Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *Journal of Clinical Investigation* 112: 1821-1830, 2003.
79. **Zimmermann R, Strauss JG, Haemmerle G, Schoiswohl G, Birner-Gruenberger R, Riederer M, Lass A, Neuberger G, Eisenhaber F, Hermetter A, and Zechner R.** Fat mobilization in adipose tissue is promoted by adipose triglyceride lipase. *Science* 306: 1383-1386, 2004.
80. **Zoladz JA, Konturek SJ, Duda K, Majerczak J, Sliwowski Z, Grandys M, and Bielanski W.** Effect of moderate incremental exercise, performed in fed and fasted state on cardio-respiratory variables and leptin and ghrelin concentrations in young healthy men. *Journal of physiology and pharmacology : an official journal of the Polish Physiological Society* 56: 63-85, 2005.

Figure legends

Figure 1. Glucose (A), insulin (C) and NEFA (E) concentrations in FASTED and FED trials and iAUC for glucose (B), insulin (D) and NEFA (F). n = 9 in both FASTED and FED trials (due to difficulty in cannulating one participant). Values are means \pm 95% CI. # denotes significant interaction effect between FASTED and FED trials by two-way ANOVA with repeated measures ($p \leq 0.05$). * denotes significantly different between FASTED versus FED trials using paired *t*-tests ($p \leq 0.05$). The shaded box in (A), (C) and (E) denotes meal time.

Figure 2. Fold changes in relative gene expression in adipose tissue under FASTED and FED trials (all n = 10, except for HSL and G0S2 n = 9 owing to Ct > 35 for one participant). The dashed line indicates baseline. Data normalised to PPIA, internal calibrator and baseline. Samples that exceeded the detectable limit (Ct > 35) were excluded from the analysis. Values are means \pm 95% CI. # denotes significant interaction effect between FASTED and FED trials using two-way ANOVA with repeated measures ($p \leq 0.05$). † denotes a time effect ($p \leq 0.05$). * comparison between baseline and 1 h post-exercise by paired *t*-tests ($p \leq 0.05$).

Figure 3. Fold changes in relative protein content in adipose tissue under FASTED and FED trials (all n = 8 due to lack of sufficient protein for two participants) (A). The dashed line indicates baseline. Data were normalised to GAPDH. Values are means \pm 95% CI. # denotes significant interaction effect between FASTED and FED trials using two-way ANOVA with repeated measures ($p \leq 0.05$). Representative images of Western blots in adipose tissue under FASTED and FED trials (B). IRS1/2, Akt2 and AMPK represents participant number 2 and 8. GLUT4, PDK4 and HSL represents participant number 3 and 4.

Figure 4. Circulating serum IL-6 (A) leptin (C) and adiponectin (E) concentrations in FASTED and FED trials (n = 9). Ex-vivo adipose tissue explant protein secretion of IL-6 (B), leptin (D) and adiponectin (F) expressed relative to L1-L4 fat mass (n = 10). Values are means \pm 95% CI. The shaded box in (A), (C) and (E) denotes meal time. † denotes a time effect from two-way ANOVA with repeated measures ($p \leq 0.05$).




Figure 1




Figure 2





Figure 3




Figure 4